

## Clinical Nursing Intervention: Allergic Dermatitis and Folliculitis<sup>\*</sup>

Kanchana Wangcharoen RN. ET.

Ruenkaew Changkittirat RN. ET.

Siriraj Hospital, Mahidol University, Thailand

### Abstract

Allergic Dermatitis and Folliculitis are peristoma skin lesion. They can make patients pain, uncomfortable, effect on their lifestyle and quality of life. And may cause of infection. However changing type of pouching system is needed. Management peristomal skin lesion with steroid<sup>1</sup> and natural substain, Manuka Honey to reduce infection, inflammation and recovery the integrity skin.<sup>2</sup>

**Keyword:** Allergic dermatitis, Folliculitis, Manuka Honey

### Introduction:

Allergic reactions causing dermatitis are less common with modern adhesives. A potential allergen provokes a response and antibody production. The inflammation of the skin usually mirrors the pattern of the allergen. A frequent allergen is the pouching system adhesive. A topical steroid preparation may reduce allergic reactions while alternative systems are tried. Folliculitis might be occurs in those who shave as removal of the pouch adhesive can cause trauma leading to folliculitis.<sup>1</sup>

### Case Report:

A 73 years old Thai elderly female was diagnosed Invasive CA Bladder S/P Open Radical cystectomy with Ileal conduit. About 4 months ago after surgery, she was pouching with Kalaya Urostomy bag. She never had skin problems. She visited the ostomy clinic because she had problems from redness skin around stoma, itching and discomfort.

---

\* Wangcharoen K, Changkittirat R. Clinical Nursing Intervention: Allergic Dermatitis and Folliculitis. In: Poster presentation of 22<sup>nd</sup> Biennial Congress, World Council of Enterostomal Therapy (WCET); 2018 Apr 14-18; Kuala Lumpur, Malaysia.





